

p -ADIC REFINABLE FUNCTIONS AND MRA-BASED WAVELETS

A. YU. KHRENNIKOV, V. M. SHELKOVICH, AND M. SKOPINA

ABSTRACT. We described a wide class of p -adic refinable equations generating p -adic multiresolution analysis. A method for the construction of p -adic orthogonal wavelet bases within the framework of the MRA theory is suggested. A realization of this method is illustrated by an example, which gives a new 3-adic wavelet basis. Another realization leads to the p -adic Haar bases which were known before.

1. INTRODUCTION

A sensation happened in the early nineties because a general scheme for the construction of wavelets was developed. This scheme is based on the notion of multiresolution analysis (MRA in the sequel) introduced by Y. Meyer and S. Mallat. Immediately specialists started to implement new wavelet systems. Nowadays it is difficult to find an engineering area where wavelets are not applied. In the p -adic setting, situation is the following. In 2002 S. V. Kozyrev [9] found a compactly supported p -adic wavelet basis for $L^2(\mathbb{Q}_p)$ which is an analog of the Haar basis. Another p -adic wavelet-type system generalizing Kozyrev's basis was constructed in [6], [7].

It turned out that the mentioned above p -adic wavelets are eigenfunctions of p -adic pseudo-differential operators [1], [2], [5]–[7], [9], [10]. This fact implies that study of wavelets is important and gives a new powerful technique for solving p -adic problems (areas of applications can be found in [4], [8], [13]).

Nevertheless, in the cited papers, *a theory describing common properties of p -adic wavelet bases and giving general methods for their construction was not developed.* Moreover, it was assumed in [3] that there are obstacles for creating a p -adic analog of the classical MRA theory. To construct a p -adic analog of a classical MRA we need a proper p -adic *refinement equation*. In [6] the following conjecture was proposed: the equality

$$(1.1) \quad \phi(x) = \sum_{r=0}^{p-1} \phi\left(\frac{1}{p}x - \frac{r}{p}\right), \quad x \in \mathbb{Q}_p,$$

may be considered as a *refinement equation*. A solution ϕ to this equation (*a refinable function*) is the characteristic function $\Omega(|x|_p)$ of the unit disc. The equation (1.1) reflects a *natural* “self-similarity” of the space $\mathbb{Q}_p$: the unit disc

2000 *Mathematics Subject Classification.* Primary 11S80, 42C40; Secondary 11E95.

Key words and phrases. p -adic multiresolution analysis; refinable equations, wavelets.

This paper was supported in part by the grant of The Swedish Royal Academy of Sciences on collaboration with scientists of former Soviet Union. The second author (V. S.) was supported in part by DFG Project 436 RUS 113/809 and Grant 05-01-04002-NNIOa of RFBR. The third author (M.S.) was supported in part by Grant 06-01-00457 of RFBR.

$B_0(0) = \{x : |x|_p \leq 1\}$ is represented as a sum of p mutually *disjoint* discs $B_0(0) = B_{-1}(0) \cup \left(\bigcup_{r=1}^{p-1} B_{-1}(r) \right)$, where $B_{-1}(r) = \{x : |x - r|_p \leq p^{-1}\}$ (see [13, I.3, Examples 1, 2.]). The equation (1.1) is an analog of the *refinement equation* generating the Haar MRA in the real analysis. Using this idea, the notion of p -adic MRA was introduced and a general scheme for its construction was described in [12]. In [12], this scheme was realized for construction 2-adic Haar MRA with using (1.1) as the generating refinement equation. In contrast to the real setting, the *refinable function* ϕ generating the Haar MRA is *periodic*, which *never holds* for real refinable functions. Due to this fact, there exist *infinity many different* orthonormal wavelet bases in the same Haar MRA. One of them coincides with Kozyrev's wavelet basis. The present paper is devoted to study of p -adic *refinement equations* generating MRAs and a method for the construction of MRA-based wavelets.

Here and in what follows, we shall use the notations and the results from [13]. Let $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{C}$ be the sets of positive integers, integers, complex numbers, respectively. The field $\mathbb{Q}_p$ of p -adic numbers is defined as the completion of the field of rational numbers $\mathbb{Q}$ with respect to the non-Archimedean p -adic norm $|\cdot|_p$. This p -adic norm is defined as follows: $|0|_p = 0$; if $x \neq 0$, $x = p^\gamma \frac{m}{n}$, where $\gamma = \gamma(x) \in \mathbb{Z}$ and the integers m, n are not divisible by p , then $|x|_p = p^{-\gamma}$. The norm $|\cdot|_p$ satisfies the strong triangle inequality $|x + y|_p \leq \max(|x|_p, |y|_p)$. The canonical form of any p -adic number $x \neq 0$ is

$$(1.2) \quad x = p^\gamma(x_0 + x_1p + x_2p^2 + \dots),$$

where $\gamma = \gamma(x) \in \mathbb{Z}$, $x_j = 0, 1, \dots, p-1$, $x_0 \neq 0$, $j = 0, 1, \dots$. We shall write the p -adic numbers $k = k_0 + k_1p + \dots + k_{s-1}p^{s-1}$, $k_0, \dots, k_{s-1} = 0, 1, 2, \dots, p-1$, in the usual (for the real analysis) form: $k = 0, 1, \dots, p^s - 1$. Denote by $B_\gamma(a) = \{x \in \mathbb{Q}_p : |x - a|_p \leq p^\gamma\}$ the disc of radius p^γ with the center at a point $a \in \mathbb{Q}_p$, $\gamma \in \mathbb{Z}$. Any two balls in $\mathbb{Q}_p$ either are disjoint or one contains the other.

There exists the Haar measure dx on $\mathbb{Q}_p$, which is positive, invariant under the shifts, i.e., $d(x + a) = dx$, and normalized by $\int_{|\xi|_p \leq 1} dx = 1$. A complex-valued function f defined on $\mathbb{Q}_p$ is called *locally-constant* if for any $x \in \mathbb{Q}_p$ there exists an integer $l(x) \in \mathbb{Z}$ such that $f(x + y) = f(x)$, $y \in B_{l(x)}(0)$. Denote by $\mathcal{D}(\mathbb{Q}_p)$ the linear spaces of locally-constant compactly supported functions (so-called test functions) [13, VI.1., 2.]. The Fourier transform of $\varphi \in \mathcal{D}(\mathbb{Q}_p)$ is defined by $\hat{\phi}(\xi) = F[\varphi](\xi) = \int_{\mathbb{Q}_p} \chi_p(\xi \cdot x) \varphi(x) dx$, $\xi \in \mathbb{Q}_p$, where $\chi_p(\xi \cdot x) = e^{2\pi i \{\xi x\}_p}$ is the additive character the field $\mathbb{Q}_p$, $\{\cdot\}_p$ is a fractional part of a number $x \in \mathbb{Q}_p$. The Fourier transform is extended to $L^2(\mathbb{Q}_p)$ in a standard way. If $f \in L^2(\mathbb{Q}_p)$, $0 \neq a \in \mathbb{Q}_p$, $b \in \mathbb{Q}_p$, then [13, VII, (3.3)]:

$$(1.3) \quad F[f(ax + b)](\xi) = |a|_p^{-1} \chi_p\left(-\frac{b}{a}\xi\right) F[f(x)]\left(\frac{\xi}{a}\right).$$

According to [13, IV, (3.1)],

$$(1.4) \quad F[\Omega(p^{-k}|\cdot|_p)](x) = p^k \Omega(p^k |x|_p), \quad k \in \mathbb{Z}, \quad x \in \mathbb{Q}_p,$$

where $\Omega(t) = 1$ for $t \in [0, 1]$; $\Omega(t) = 0$ for $t \notin [0, 1]$.

2. MULTIREOLUTION ANALYSIS

Let $I_p = \{a = p^{-\gamma}(a_0 + a_1p + \cdots + a_{\gamma-1}p^{\gamma-1}) : \gamma \in \mathbb{N}; a_j = 0, 1, \dots, p-1; j = 0, 1, \dots, \gamma-1\}$. It is well known that $\mathbb{Q}_p = B_0(0) \cup \bigcup_{\gamma=1}^{\infty} S_\gamma$, where $S_\gamma = \{x \in \mathbb{Q}_p : |x|_p = p^{-\gamma}\}$. Due to (1.2), $x \in S_\gamma$, $\gamma \geq 1$, if and only if $x = x_{-\gamma}p^{-\gamma} + x_{-\gamma+1}p^{-\gamma+1} + \cdots + x_{-1}p^{-1} + \xi$, where $x_{-\gamma} \neq 0$, $\xi \in B_0(0)$. Since $x_{-\gamma}p^{-\gamma} + x_{-\gamma+1}p^{-\gamma+1} + \cdots + x_{-1}p^{-1} \in I_p$, we have a “natural” decomposition of $\mathbb{Q}_p$ to a union of mutually disjoint discs: $\mathbb{Q}_p = \bigcup_{a \in I_p} B_0(a)$. So, I_p is a “natural” set of shifts for $\mathbb{Q}_p$.

Definition 2.1. ([12]) A collection of closed spaces $V_j \subset L^2(\mathbb{Q}_p)$, $j \in \mathbb{Z}$, is called a *multiresolution analysis (MRA) in $L^2(\mathbb{Q}_p)$* if the following axioms hold

- (a) $V_j \subset V_{j+1}$ for all $j \in \mathbb{Z}$;
- (b) $\bigcup_{j \in \mathbb{Z}} V_j$ is dense in $L^2(\mathbb{Q}_p)$;
- (c) $\bigcap_{j \in \mathbb{Z}} V_j = \{0\}$;
- (d) $f(\cdot) \in V_j \iff f(p^{-1}\cdot) \in V_{j+1}$ for all $j \in \mathbb{Z}$;
- (e) there exists a function $\phi \in V_0$ such that the system $\{\phi(\cdot - a), a \in I_p\}$ is an orthonormal basis for V_0 .

The function ϕ from axiom (e) is called *scaling*. It follows immediately from axioms (d) and (e) that the functions $p^{j/2}\phi(p^{-j}\cdot - a)$, $a \in I_p$, form an orthonormal basis for V_j , $j \in \mathbb{Z}$. According to the standard scheme (see, e.g., [11, §1.3]) for construction of MRA-based wavelets, for each j , we define a space W_j (*wavelet space*) as the orthogonal complement of V_j in V_{j+1} , i.e., $V_{j+1} = V_j \oplus W_j$, $j \in \mathbb{Z}$, where $W_j \perp V_j$, $j \in \mathbb{Z}$. It is not difficult to see that

$$(2.1) \quad f \in W_j \iff f(p^{-1}\cdot) \in W_{j+1}, \quad \text{for all } j \in \mathbb{Z}$$

and $W_j \perp W_k$, $j \neq k$. Taking into account axioms (b) and (c), we obtain

$$(2.2) \quad \bigoplus_{j \in \mathbb{Z}} W_j = L^2(\mathbb{Q}_p) \quad (\text{orthogonal direct sum}).$$

If now we find a finite number of functions $\psi^{(\nu)} \in W_0$, $\nu \in A$, such that the system $\{\psi^{(\nu)}(x - a), a \in I_p, \nu \in A\}$ is an orthonormal basis for W_0 , then, due to (2.1) and (2.2), the system $\{p^{j/2}\psi^{(\nu)}(p^{-j}\cdot - a), a \in I_p, j \in \mathbb{Z}, \nu \in A\}$, is an orthonormal basis for $L^2(\mathbb{Q}_p)$. Such functions $\psi^{(\nu)}$ are called *wavelet functions* and the basis is a *wavelet basis*.

Let ϕ be a *scaling function* for a MRA. As was mentioned above, the system $\{p^{1/2}\phi(p^{-1}\cdot - a), a \in I_p\}$ is a basis for V_1 . It follows from axiom (a) that

$$(2.3) \quad \phi = \sum_{a \in I_p} \beta_a \phi(p^{-1}\cdot - a), \quad \beta_a \in \mathbb{C}.$$

We see that the function ϕ is a solution of a special kind of functional equation. Such equations are called *refinement equations*, and their solutions are called *refinable functions*¹.

A natural way for the construction of a MRA (see, e.g., [11, §1.2]) is the following. We start with an appropriate function ϕ whose I_p -shifts form an orthonormal system and set

$$(2.4) \quad V_j = \overline{\text{span}\{\phi(p^{-j}\cdot - a) : a \in I_p\}}, \quad j \in \mathbb{Z}.$$

¹Usually the terms “scaling function” and “refinable function” are synonyms in the literature, and they are used for both the senses: as a solution to a refinement equation and as a function generating MRA. We separate the meanings of these terms.

It is clear that axioms (d) and (e) of Definition 2.1 are fulfilled. Of course, not any such a function ϕ provides axiom (a). In the *real setting*, the relation $V_0 \subset V_1$ holds if and only if the scaling function satisfies a refinement equation. Situation is different in p -adics. Generally speaking, a refinement equation (2.3) *does not imply the including property* $V_0 \subset V_1$ because the set of the shifts I_p *does not form a group*. Indeed, we need all the functions $\phi(\cdot - b)$, $b \in I_p$, to belong to the space V_1 , i.e., the identities $\phi(x - b) = \sum_{a \in I_p} \alpha_{a,b} \phi(p^{-1}x - a)$ should be fulfilled for all $b \in I_p$. Since $p^{-1}b + a$ is not in I_p in general, we *can not state* that $\phi(x - b) = \sum_{a \in I_p} \alpha_{a,b} \phi(p^{-1}x - p^{-1}b - a) \in V_1$ for all $b \in I_p$. Nevertheless, some refinement equations imply including property, which may happen because of different causes.

The *refinement equation* (1.1) is a particular case of (2.3).

3. CONSTRUCTION OF REFINABLE FUNCTIONS

Now we are going to study p -adic refinement equations and their solutions. We restrict ourselves by the refinement equations (2.3) with a finite number of the terms in the right-hand side:

$$(3.1) \quad \phi(x) = \sum_{k=0}^{p^s-1} \beta_k \phi\left(\frac{1}{p}x - \frac{k}{p^s}\right).$$

If $\phi \in L^2(\mathbb{Q}_p)$, taking the Fourier transform and using (1.3), one can rewrite (3.1) as

$$(3.2) \quad \widehat{\phi}(\xi) = m_0\left(\frac{\xi}{p^{s-1}}\right) \widehat{\phi}(p\xi),$$

where

$$(3.3) \quad m_0(\xi) = \frac{1}{p} \sum_{k=0}^{p^s-1} \beta_k \chi_p(k\xi)$$

is a trigonometric polynomial. It is clear that $m_0(0) = 1$ whenever $\widehat{\phi}(0) \neq 0$.

Theorem 3.1. *If ϕ is a refinable function such that $\text{supp } \widehat{\phi} \subset B_0(0)$ and the system $\{\phi(x - a) : a \in I_p\}$ is orthonormal, then axiom (a) from Definition 2.1 holds for the spaces (2.4).*

Proof. Since $\chi_p(-\xi) = 1$ for $\xi \in B_0(0)$, we have $\chi_p(-\xi) \widehat{\phi}(\xi) = \widehat{\phi}(\xi)$. Applying the Fourier transform, we obtain $\phi(x + 1) = \phi(x)$. Thus ϕ is a 1-*periodic* function. Since for all $a \in I_p$ and all $k = 0, 1, \dots, p^s - 1$, either $\frac{a}{p} + \frac{k}{p^s} \in I_p$, or $\frac{a}{p} + \frac{k}{p^s} - 1 \in I_p$. Due to the 1-*periodicity* of ϕ , it follows from (3.1) that $\phi(x - b) \in V_1$ for all $b \in I_p$. This implies $V_0 \subset V_1$, similarly $V_j \subset V_{j+1}$ for any $j \in \mathbb{Z}$. $\square$

Theorem 3.2. *Let $\phi \in L^2(\mathbb{Q}_p)$ be a refinable function, the system $\{\phi(x - a) : a \in I_p\}$ be orthonormal and $\text{supp } \widehat{\phi} \subset B_0(0)$. Axiom (b) of Definition 2.1 holds for the spaces (2.4) (i.e., $\overline{\cup_{j \in \mathbb{Z}} V_j} = L^2(\mathbb{Q}_p)$) if and only if*

$$(3.4) \quad \bigcup_{j \in \mathbb{Z}} \text{supp } \widehat{\phi}(p^j \cdot) = \mathbb{Q}_p.$$

Proof. First of all we note that, due to axioms (d) and (e), each space V_j is invariant with respect to the shifts $t = p^j a$, $a \in I_p$. Show that the space $\overline{\cup_{j \in \mathbb{Z}} V_j}$ is invariant with respect to any shift $t \in \mathbb{Q}_p$. Every $t \in \mathbb{Q}_p$ may be approximated by a vector $p^j a$, $a \in I_p$, with arbitrary large $j \in \mathbb{Z}$. If $f \in \cup_{j \in \mathbb{Z}} V_j$, by axiom (a) (which holds due to Theorem 3.1), then $f \in V_j$ for all $j \geq j_1$. It follows from the continuity of the function $\|f(\cdot + t)\|_2$ that $f(\cdot + t) \in \overline{\cup_{j \in \mathbb{Z}} V_j}$. Now let $t \in \mathbb{Q}_p$. If $g \in \overline{\cup_{j \in \mathbb{Z}} V_j}$, then approximating g by the functions $f \in \cup_{j \in \mathbb{Z}} V_j$, again using the continuity of the shift operator and the invariance of L_2 norm with respect to the shifts, we derive $g(\cdot + t) \in \overline{\cup_{j \in \mathbb{Z}} V_j}$. For $X \subset L^2(\mathbb{Q}_p)$, set $\hat{X} = \{\hat{f} : f \in X\}$. By the Wiener theorem for L_2 (see, e.j., [11], all the arguments of the proof given there may be repeated word for word with replacing $\mathbb{R}$ by $\mathbb{Q}_p$), a closed subspace X of the space $L^2(\mathbb{Q}_p)$ is invariant with respect to the shifts if and only if $\hat{X} = L_2(\Omega)$ for some set $\Omega \subset \mathbb{Q}_p$. Let $X = \overline{\cup_{j \in \mathbb{Z}} V_j}$, then $\hat{X} = L_2(\Omega)$. Thus $X = L^2(\mathbb{Q}_p)$ if and only if $\Omega = \mathbb{Q}_p$. Set $\phi_j = \phi(p^{-j} \cdot)$, $\Omega_0 = \cup_{j \in \mathbb{Z}} \text{supp } \hat{\phi}_j$ and prove that $\Omega = \Omega_0$. Since $\phi_j \in V_j$, $j \in \mathbb{Z}$, we have $\text{supp } \hat{\phi}_j \subset \Omega$, and hence $\Omega_0 \subset \Omega$. Now assume that $\Omega \setminus \Omega_0$ contains a set of positive measure Ω_1 . If $f \in V_j$, taking the Fourier transform from the expansion $f = \sum_{a \in I_p} h_a \phi(p^{-j} \cdot - a)$ we see that $\hat{f} = 0$ almost everywhere on Ω_1 . Hence the same is true for any $f \in \cup_{j \in \mathbb{Z}} V_j$. Passing to the limit we deduce that the Fourier transform of any $f \in X$ is equal to zero almost everywhere on Ω_1 , i.e., $L_2(\Omega) = L_2(\Omega_0)$. It remains to note that $\text{supp } \hat{\phi}_j = \text{supp } \hat{\phi}(p^j \cdot)$ $\square$

Theorem 3.3. *If $\phi \in L^2(\mathbb{Q}_p)$ and the system $\{\phi(x - a) : a \in I_p\}$ is orthonormal, then axiom (c) of Definition 2.1 holds, i.e., $\cap_{j \in \mathbb{Z}} V_j = \{0\}$.*

Proof. First, using the standard scheme (see, e.g., [11, Lemma 1.2.8.]), we prove that for any $f \in L^2(\mathbb{Q}_p)$

$$(3.5) \quad \lim_{j \rightarrow -\infty} \sum_{a \in I_p} |(f, p^{j/2} \phi(p^{-j} \cdot - a))|^2 = 0,$$

where $(\cdot, \cdot)$ is the scalar product in $L^2(\mathbb{Q}_p)$. Since the space $\mathcal{D}(\mathbb{Q}_p)$ is dense in $L^2(\mathbb{Q}_p)$ [13, VI.2], it suffices to prove (3.5) for any $\varphi \in \mathcal{D}(\mathbb{Q}_p)$. If $\varphi \in \mathcal{D}(\mathbb{Q}_p)$, then there exists N such that $\varphi(x) = 0$ for all $|\xi|_p > p^N$. Since $|\varphi(x)| \leq M$ for all $|\xi|_p \leq p^N$, we have

$$\begin{aligned} \sum_{a \in I_p} |(\varphi(\cdot), p^{j/2} \phi(p^{-j} \cdot - a))|^2 &\leq p^j \sum_{a \in I_p} \left(\int_{|x|_p \leq p^N} |\varphi(x)| |\phi(p^{-j} x - a)| dx \right)^2 \\ &\leq p^{j+N} M^2 \sum_{a \in I_p} \int_{|x|_p \leq p^N} |\phi(p^{-j} x - a)|^2 dx. \end{aligned}$$

By the change of variables $\eta = p^{-j} x - a$, we obtain

$$\sum_{a \in I_p} |(\varphi(\cdot), p^{j/2} \phi(p^{-j} \cdot - a))|^2 \leq M^2 p^N \int_{A_{Nj}} |\phi(\eta)|^2 d\eta = M^2 p^N \int_{\mathbb{Q}_p} \theta_{Nj}(\eta) |\phi(\eta)|^2 d\eta,$$

where θ_{Nj} is the characteristic function of the set $A_{Nj} = \cup_{a \in I_p} \{\eta : |\eta + a|_p \leq 2^{N+j}\}$. Since $\lim_{j \rightarrow -\infty} \theta_{Nj}(\eta) = 0$ for any $\eta \neq -a$, using the Lebesgue dominated convergence theorem [13, IV.4], we obtain $\lim_{j \rightarrow -\infty} \int_{\mathbb{Q}_p} \theta_{Nj}(\eta) |\phi(\eta)|^2 d\eta = 0$.

If now we assume that $f \in \cap_{j \in \mathbb{Z}} V_j$, then $f \in V_j$ for all $j \in \mathbb{Z}$ and, due to (3.5),

$$\|f\| = \left(\sum_{a \in I_p} |(f, p^{j/2} \phi(p^{-j} \cdot -a))|^2 \right)^{1/2} \rightarrow 0, \quad j \rightarrow -\infty,$$

i.e., $\|f\| = 0$ which was to be proved. $\square$

Theorem 3.4. *Let ϕ be a refinable function such that $\text{supp } \widehat{\phi} \subset B_0(0)$. If $|\widehat{\phi}(\xi)| = 1$ for all $\xi \in B_0(0)$ then the system $\{\phi(x-a) : a \in I_p\}$ is orthonormal.*

Proof. Taking into account formula (1.4), using the inclusion $\text{supp } \widehat{\phi} \subset B_0(0)$ and the Plancherel formula, we have for any $a \in I_p$

$$\begin{aligned} (\phi(\cdot), \phi(\cdot - a)) &= \int_{\mathbb{Q}_p} \phi(x) \overline{\phi}(x-a) dx = \int_{B_0(0)} |\widehat{\phi}(\xi)|^2 \chi_p(a\xi) d\xi \\ &= \int_{B_0(0)} \chi_p(a\xi) d\xi = \int_{\mathbb{Q}_p} \Omega(|\xi|_p) \chi_p(a\xi) d\xi = \Omega(|a|_p) = \delta_{a0}. \end{aligned}$$

$\square$

So, to construct a MRA we can take a function ϕ for which the hypotheses of Theorems 3.4 and (3.4) are fulfilled. Next we are going to describe all such functions.

Proposition 3.5. *If $\phi \in L^2(\mathbb{Q}_p)$ is a solution of refinable equation (3.2), $\widehat{\phi}(\xi)$ is continuous at the point 0 and $\widehat{\phi}(0) \neq 0$, then*

$$(3.6) \quad \widehat{\phi}(\xi) = \widehat{\phi}(0) \prod_{j=1}^{\infty} m_0\left(\frac{\xi}{p^{s-j}}\right).$$

Proof. Iterating (3.2) N times, $N \geq 1$, we have

$$(3.7) \quad \widehat{\phi}(\xi) = \prod_{j=1}^N m_0\left(\frac{\xi}{p^{s-j}}\right) \widehat{\phi}(p^N \xi).$$

Taking into account that $\widehat{\phi}(\xi)$ is continuous at the point 0 and the fact that $|p^N \xi|_p = p^{-N} |\xi|_p \rightarrow 0$ as $N \rightarrow +\infty$ for any $\xi \in \mathbb{Q}_p$, we obtain (3.6). $\square$

Proposition 3.6. *If $\widehat{\phi}$ is defined by (3.6), where m_0 is a trigonometric polynomial (3.3), $m_0(0) = 1$, then (3.2) holds. Furthermore, if $\xi \in \mathbb{Q}_p$ such that $|\xi|_p = p^{-n}$, then $\widehat{\phi}(\xi) = \widehat{\phi}(0)$ for $n \geq s-1$, and*

$$(3.8) \quad \widehat{\phi}(\xi) = \widehat{\phi}(0) \prod_{j=1}^{s-n-1} m_0\left(\frac{\xi}{p^{s-j}}\right) = \widehat{\phi}(0) \prod_{j=1}^{s-n-1} m_0\left(\frac{\tilde{\xi}}{p^{s-j}}\right) = \widehat{\phi}(\tilde{\xi})$$

where $\tilde{\xi} = \xi_n p^n + \xi_{n+1} p^{n+1} + \dots + \xi_{s-2} p^{s-2}$, $\xi_n \neq 0$, for $n \leq s-2$.

Proof. Relation (3.6) implies $\widehat{\phi}(p\xi) = \widehat{\phi}(0) \prod_{j=1}^{\infty} m_0\left(\frac{\xi}{p^{s-j-1}}\right)$ and, consequently, (3.2). Let $n \leq s-2$, $|\xi|_p = p^{-n}$, i.e., $\xi = \xi_n p^n + \xi_1 p^{n+1} + \dots$, $\xi_n \neq 0$. Since $\chi_p\left(\frac{k\xi'}{p^{s-j}}\right) = 1$, whenever $\xi' \in B_{-s+1}(0)$, $j \in \mathbb{N}$, $k = 0, 1, \dots, p^s - 1$, and $\chi_p\left(\frac{k\xi}{p^{s-j}}\right) = 1$, whenever $j \geq s-n$, we have (3.8). It is clear that $\widehat{\phi}(\xi) = \widehat{\phi}(0)$ for $n \geq s-1$. $\square$

Corollary 3.7. *The function $\widehat{\phi}$ from Proposition 3.6 is locally-constant. Moreover, if $M \geq -s + 2$, $\text{supp} \widehat{\phi} \subset B_M(0)$, then for any $k = 0, 1, \dots, p^{M+s-1} - 1$ and for all $x \in B_{s-1}(\frac{k}{p^M})$ we have $\widehat{\phi}(x) = \widehat{\phi}(\frac{k}{p^M})$.*

Proposition 3.8. *Let $\widehat{\phi}$ be defined by (3.6), where m_0 is a trigonometric polynomial (3.3). If $m_0(0) = 1$, $m_0(\frac{k}{p^s}) = 0$ for all $k = 1, \dots, p^s - 1$ which are not divisible by p , then $\text{supp} \widehat{\phi} \subset B_0(0)$, $\widehat{\phi} \in L^2(\mathbb{Q}_p)$. If, furthermore, $|m_0(\frac{k}{p^s})| = 1$ for all $k = 1, \dots, p^s - 1$ which are divisible by p , then $|\widehat{\phi}(x)| = |\widehat{\phi}(0)|$ for any $x \in B_0(0)$.*

Proof. By Proposition 3.6, $\widehat{\phi}$ satisfies (3.2) and (3.8). Let us check that $\widehat{\phi}(\xi) = 0$ for all ξ such that $|\xi|_p = p^M$, $M \geq 1$. It follows from (3.2) that it suffices to consider only $M = 1$. Let $|\xi|_p = p$, i.e., $\xi = \frac{1}{p}\xi_{-1} + \xi_0 + \xi_1 p + \dots + \xi_{s-2} p^{s-2} + \xi'$, where $\xi_{-1} \neq 0$, $\xi' \in B_{-s+1}(0)$. In view of (3.8), we have $\widehat{\phi}(\xi) = \widehat{\phi}(\xi) = \widehat{\phi}(\frac{k}{p})$, where $k = \xi_{-1} + \xi_0 p + \xi_1 p^2 + \dots + \xi_{s-2} p^{s-1}$, $\xi_{-1} \neq 0$. Note that the first factor of the product in (3.8) is $m_0(\frac{k}{p^s}) = 0$. Thus $\widehat{\phi}(\xi) = 0$ for all ξ such that $|\xi|_p = p$. The rest statements follow from Propositions 3.5, 3.6. $\square$

Due to Theorems 3.1-3.4, the refinable functions with masks satisfying the hypotheses of Proposition 3.8 generate MRAs. Next, we will see that all properties of a mask m_0 described in Propositions 3.8 are necessary for the corresponding refinable function ϕ to be such that $\text{supp} \widehat{\phi} \subset B_0(0)$ and the system $\{\phi(x - a) : a \in I_p\}$ is orthonormal.

Theorem 3.9. *Let $\widehat{\phi}$ be defined by (3.6), where m_0 is a trigonometric polynomial (3.3). If $\text{supp} \widehat{\phi} \subset B_0(0)$ and the system $\{\phi(x - a) : a \in I_p\}$ is orthonormal, then $|m_0(\frac{k}{p^s})| = 0$ whenever k is not divisible by p , and $|m_0(\frac{k}{p^s})| = 1$ whenever k is divisible by p , $k = 1, 2, \dots, p^s - 1$.*

Proof. Let $a \in I_p$. Due to the orthonormality of $\{\phi(x - a) : a \in I_p\}$, using the Plancherel formula and Corollary 3.7, we have

$$\begin{aligned} \delta_{a0} &= (\phi(\cdot), \phi(\cdot - a)) = \int_{\mathbb{Q}_p} \phi(x) \overline{\phi}(x - a) dx = \int_{B_0(0)} |\widehat{\phi}(\xi)|^2 \chi_p(a\xi) d\xi \\ &= \sum_{k=0}^{p^{s-1}-1} \int_{|\xi-k|_p \leq p^{-s+1}} |\widehat{\phi}(\xi)|^2 \chi_p(a\xi) d\xi = \sum_{k=0}^{p^{s-1}-1} |\widehat{\phi}(k)|^2 \int_{|\xi-k|_p \leq p^{-s+1}} \chi_p(a\xi) d\xi \\ &= \sum_{k=0}^{p^{s-1}-1} |\widehat{\phi}(k)|^2 \chi_p(ak) \int_{|\xi|_p \leq p^{-s+1}} \chi_p(a\xi) d\xi = \frac{1}{p^{s-1}} \Omega(|p^{s-1}a|_p) \sum_{k=0}^{p^{s-1}-1} |\widehat{\phi}(k)|^2 \chi_p(ak). \end{aligned}$$

Since $\Omega(|p^{s-1}a|_p) \neq 0$ for all $a \in B_{-s+1}(0)$, this yields

$$\frac{1}{p^{s-1}} \sum_{k=0}^{p^{s-1}-1} |\widehat{\phi}(k)|^2 \chi_p(ak) = \delta_{a0}, \quad a = 0, \frac{1}{p^{s-1}}, \dots, \frac{p^{s-1}-1}{p^{s-1}}.$$

Consider these equalities as a linear system with respect to the unknowns $z_k = |\widehat{\phi}(k)|^2$. It is well-known that the system has a unique solution $z_k = 1$, $k =$

$0, 1, \dots, p^{s-1} - 1$, i.e., $|\widehat{\phi}(k)| = 1$ for all $k = 0, 1, \dots, p^{s-1} - 1$. In particular, it follows that $|\widehat{\phi}(0)|^2 = 1$, which reduces (3.6) to

$$(3.9) \quad \widehat{\phi}(\xi) = m_0\left(\frac{\xi}{p^{s-1}}\right)m_0\left(\frac{\xi}{p^{s-2}}\right)\cdots m_0(\xi).$$

Let us check that $|m_0(\frac{k}{p^s})| = 1$ for all k divisible by p , $k = 1, 2, \dots, p^{s-1} - 1$. This is equivalent to $|m_0(\frac{k}{p^N})| = 1$ whenever $N = 1, 2, \dots, p^{s-1}$, $k = 1, 2, \dots, p^N - 1$, k is not divisible by p . We will prove this statement by induction on N . For the inductive base with $N = 1$, note that $1 = |\widehat{\phi}(p^{s-2})| = |m_0(\frac{1}{p})|$. For the inductive step, assume that $|m_0(\frac{k}{p^n})| = 1$ for all $n = 1, 2, \dots, N$, $N \leq s - 2$, $k = 1, 2, \dots, p^n - 1$, k is not divisible by p . Using (3.9) and 1-periodicity of m_0 , we have $1 = |\widehat{\phi}(lp^{s-N-2})| = |m_0(\frac{l}{p^{N+1}})m_0(\frac{l}{p^N})\cdots m_0(\frac{l}{p^{N-s+2}})| = |m_0(\frac{l}{p^{N+1}})|$, for all $l = 1, 2, \dots, p^{N+1} - 1$, l is not divisible by p . Now assume that $m_0(\frac{k}{p^s}) \neq 0$ for some $k = 1, 2, \dots, p^s - 1$ not divisible by p . Since $\widehat{\phi}(\frac{k}{p}) = 0$, due to (3.9), there exists $n = 1, 2, \dots, s$ such that $m_0(\frac{k}{p^{s-n}}) = 0$ which contradicts to $|m_0(\frac{k}{p^{s-n}})| = 1$. $\square$

We have investigated refinable functions whose Fourier transform is supported in the unit disk $B_0(0)$. Such functions provide axiom (a) of Definition 2.1 because of a trivial argument given in Theorem 3.1. If $\text{supp } \widehat{\phi} \not\subset B_0(0)$, generally speaking, the relation $\phi(\cdot - a) \in V_1$ does not follow from the refinability of ϕ for all $a \in I_p$ because I_p is not a group. Nevertheless, we observed that some such refinable functions also provide axiom (a). Let $p = 2$, $s = 3$, ϕ be defined by (3.6), where m_0 is given by (3.3), $m_0(1/4) = m_0(3/8) = m_0(7/16) = m_0(15/16) = 0$. It is not difficult to see that $\text{supp } \widehat{\phi} \subset B_1(0)$, $\text{supp } \widehat{\phi} \not\subset B_0(0)$. Evidently, axiom (a) will be fulfilled whenever $\phi(x - \frac{k}{4}) = \sum_{r=0}^7 \gamma_{kr} \phi(\frac{1}{2}x - \frac{r}{8})$, $k = 1, 2, 3$, $x \in \mathbb{Q}_2$, which is equivalent to $\widehat{\phi}(\xi)\chi_2(\frac{k\xi}{4}) = m_k(\frac{\xi}{4})\widehat{\phi}(2\xi)$, $k = 1, 2, 3$, $\xi \in \mathbb{Q}_2$, where $m_k(\xi) = \frac{1}{2} \sum_{r=0}^7 \gamma_{k,r} \chi_2(r\xi)$. Combining this with (3.2) we have $\widehat{\phi}(8\xi)(m_0(\xi)\chi_2(k\xi) - m_k(\xi)) = 0$, $k = 1, 2, 3$, $\xi \in \mathbb{Q}_2$. These equalities will be fulfilled for any $\xi \in \mathbb{Q}_2$ whenever they are fulfilled for $\xi = l/16$, $l = 0, 1, \dots, 15$. Desirable polynomials m_k , $k = 1, 2, 3$, exist because we have $\widehat{\phi}(\frac{1}{2}) = \widehat{\phi}(\frac{3}{2}) = \widehat{\phi}(\frac{5}{2}) = \widehat{\phi}(\frac{9}{2}) = \widehat{\phi}(\frac{11}{2}) = \widehat{\phi}(\frac{13}{2}) = \widehat{\phi}(1) = \widehat{\phi}(5) = 0$. So, we succeeded with providing axiom (a) of Definition 2.1, but, unfortunately, such a ϕ is not a scaling function generating MRA because axiom (e) is not valid. Moreover, it is possible to show that for any refinable function whose Fourier transform is in $B_1(0)$ but not in $B_0(0)$ the shift system $\{\phi(x - a) : a \in I_p\}$ is not orthogonal. We suggest the following conjecture: *it does not exist compactly supported refinable functions with mutually orthogonal shifts $\{\phi(x - a) : a \in I_p\}$ whose Fourier transform is not supported in $B_0(0)$.*

4. WAVELET BASES

Now we discuss how to find wavelet functions if we have already a p -adic MRA generating by scaling function. Let the refinement equation for ϕ be (3.1). We look for wavelet functions $\psi^{(\nu)}$, $\nu = 1, \dots, p-1$, in the form $\psi^{(\nu)}(x) = \sum_{k=0}^{p^s-1} \gamma_{\nu k} \phi(\frac{1}{p}x -$

$\frac{k}{p^s})$, where the coefficients $\gamma_{\nu k}$ are chosen such that

$$(4.1) \quad (\psi^{(\nu)}, \phi(\cdot - a)) = 0, \quad (\psi^{(\nu)}, \psi^{(\mu)}(\cdot - a)) = \delta_{\nu\mu} \delta_{0a}, \quad \nu, \mu = 1, \dots, p-1,$$

for any $a \in I_p$. It is clear that (4.1) are fulfilled for all $a \neq 0, \frac{1}{p^{s-1}}, \dots, \frac{p^{s-1}-1}{p^{s-1}}$.

Set $B = \frac{1}{\sqrt{p}}(\beta_0, \dots, \beta_{p^s-1})^T$, $G_\nu = \frac{1}{\sqrt{p}}(\gamma_{\nu 0}, \dots, \gamma_{\nu, p^s-1})^T$, $\nu = 1, \dots, p-1$,

$$S = \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & 0 \end{pmatrix}.$$

To provide (4.1) for $a = 0, \frac{1}{p^{s-1}}, \dots, \frac{p^{s-1}-1}{p^{s-1}}$ we should find vectors $G_1, \dots, G_{p-1}$ so that the matrix

$$U = (S^0 B, \dots, S^{p^{s-1}-1} B, S^0 G_1, \dots, S^{p^{s-1}-1} G_1, \dots, S^0 G_{p-1}, \dots, S^{p^{s-1}-1} G_{p-1})$$

is unitary.

Example 4.1. Let $s = 1$. According to Proposition 3.8, we set $m_0(0) = 1$, $m_0(\frac{k}{p}) = 0$, $k = 1, \dots, p-1$ and find the mask $m_0(\xi) = \frac{1}{p} \sum_{k=0}^{p-1} \chi_p(k\xi)$ (here $\beta_k = 1$ in (3.3) and $B = \frac{1}{\sqrt{p}}(1, \dots, 1)^T$). The corresponding refinement equation (3.1) coincides with the “natural” refinement equation (1.1), and its solution $\Omega(| \cdot |_p)$ is a refinable function generating a MRA because of Theorems 3.1-3.4. To find wavelets we observe that the unitary matrix $\{\frac{1}{\sqrt{p}} e^{2\pi i k l}\}_{k,l=0,\dots,p-1}$ may be taken as U^T . Computing the wavelet functions corresponding to this matrix U , we derive the formulas which were found in [9]: $\psi^{(\nu)}(x) = \chi_p(\frac{\nu}{p}x) \Omega(|x|_p)$, $\nu = 1, \dots, p-1$.

Example 4.2. Let $s = 2$, $p = 3$. According to Proposition 3.8, we set $m_0(\frac{k}{9}) = 0$ if k is not divisible by 3, and $m_0(0) = 1$, $m_0(\frac{1}{3}) = m_0(\frac{2}{3}) = -1$. In this case $m_0(z) = 3^{-2}(-1 + 2z + 2z^2 - z^3 + 2z^4 + 2z^5 - z^6 + 2z^7 + 2z^8)$, $z = e^{2\pi i \xi}$ and

$$\widehat{\phi}(\xi) = \begin{cases} 1, & |\xi|_p \leq \frac{1}{3}, \\ -1, & |\xi - 1|_p \leq \frac{1}{3}, \\ -1, & |\xi - 2|_p \leq \frac{1}{3}, \\ 0, & |\xi|_p \geq 3. \end{cases}$$

Thus, the corresponding refinement equation (3.1) is $\phi(x) = \sum_{k=0}^8 \beta_k \phi(\frac{x}{3} - \frac{k}{9})$, where $\beta_0 = -\frac{1}{3}$, $\beta_1 = \beta_2 = \frac{2}{3}$, $\beta_3 = -\frac{1}{3}$, $\beta_4 = \beta_5 = \frac{2}{3}$, $\beta_6 = -\frac{1}{3}$, $\beta_7 = \beta_8 = \frac{2}{3}$, and

$$(4.2) \quad \phi(x) = \begin{cases} -\frac{1}{3}, & |x|_p \leq \frac{1}{3} \\ \frac{2}{3}, & |x - \frac{1}{3}|_p \leq 1 \\ \frac{2}{3}, & |x - \frac{2}{3}|_p \leq 1 \\ 0, & |x|_p \geq 9 \end{cases} = \frac{1}{3} \Omega(|3x|_p) (1 - e^{2\pi i \{x\}_3} - e^{4\pi i \{x\}_3}).$$

So, we have $B = \frac{1}{3\sqrt{3}}(-1, 2, 2, -1, 2, 2, -1, 2, 2)^T$, and the vectors B, SB, S^2B are orthonormal. We need to extend these three columns to a unitary matrix $U = (B, SB, S^2B, G_1, SG_1, S^2G_1, G_2, SG_2, S^2G_2)$. It is not difficult to see that the vectors $G_1 = \frac{1}{\sqrt{3}}(1, 0, 0, -1, 0, 0, 0, 0, 0)^T$, $G_2 = \frac{1}{\sqrt{3}}(1, 0, 0, 1, 0, 0, -2, 0, 0)^T$ are

appropriate. The corresponding wavelet functions are $\psi^{(1)} = \sqrt{\frac{3}{2}}(\phi(\frac{x}{3}) - \phi(\frac{x}{3} - \frac{1}{3}))$, $\psi^{(2)} = \frac{1}{\sqrt{2}}(\phi(\frac{x}{3}) + \phi(\frac{x}{3} - \frac{1}{3}) - 2\phi(\frac{x}{3} - \frac{2}{3}))$. Substituting (4.2), we obtain

$$\psi^{(1)}(x) = \begin{cases} -\sqrt{\frac{3}{2}}, & |x|_p \leq \frac{1}{3}, \\ \sqrt{\frac{3}{2}}, & |x - 1|_p \leq \frac{1}{3}, \\ 0, & |x - 2|_p \leq \frac{1}{3}, \\ 0, & |x|_p \geq 3; \end{cases} \quad \psi^{(2)}(x) = \begin{cases} -\frac{1}{\sqrt{2}}, & |x|_p \leq \frac{1}{3}, \\ -\frac{1}{\sqrt{2}}, & |x - 1|_p \leq \frac{1}{3}, \\ \sqrt{2}, & |x - 2|_p \leq \frac{1}{3}, \\ 0, & |x|_p \geq 3. \end{cases}$$

REFERENCES

1. S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, *Harmonic analysis in the p-adic Lizorkin spaces: fractional operators, pseudo-differential equations, p-adic wavelets, Tauberian theorems*, Journal of Fourier Analysis and Applications, Vol. 12, Issue 4 (2006), 393–425.
2. S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, *p-Adic semi-linear evolutionary pseudo-differential equations in the Lizorkin space*, Dokl. Ross. Akad. Nauk, **415**, no. 3, (2007), 295–299. English transl. in Russian Doklady Mathematics, **76**, no. 1, (2007), 539–543.
3. J.J. Benedetto, and R.L. Benedetto, *A wavelet theory for local fields and related groups*, The Journal of Geometric Analysis **3** (2004), 423–456.
4. A. Khrennikov, *Non-archimedean analysis: quantum paradoxes, dynamical systems and biological models*. Kluwer Academic Publ., Dordrecht, 1997.
5. A.Yu. Khrennikov, and S.V. Kozyrev, *Pseudodifferential operators on ultrametric spaces and ultrametric wavelets*, Izvestia Akademii Nauk, Seria Math. **69** no. 5 (2005), 133–148.
6. A.Yu. Khrennikov, V.M. Shelkovich, *p-Adic multidimensional wavelets and their application to p-adic pseudo-differential operators*. <http://arxiv.org/abs/math-ph/0612049>
7. A.Yu. Khrennikov, V.M. Shelkovich, *Non-Haar p-adic wavelets and pseudo-differential operators*, Dokl. Ross. Akad. Nauk, **418**, no. 2, (2008). English transl. in Russian Doklady Mathematics, (2008).
8. A.N. Kochubei, *Pseudo-differential equations and stochastics over non-archimedean fields*, Marcel Dekker. Inc. New York, Basel, 2001.
9. S.V. Kozyrev, *Wavelet analysis as a p-adic spectral analysis*, Izvestia Akademii Nauk, Seria Math. **66** no. 2 (2002), 149–158.
10. S.V. Kozyrev, *p-Adic pseudodifferential operators and p-adic wavelets*, Theor. Math. Physics **138**, no. 3 (2004), 1–42.
11. I. Novikov, V. Protassov, and M. Skopina, *Wavelet Theory*. Moscow: Fizmatlit, 2005.
12. V. M. Shelkovich, M. Skopina *p-Adic Haar multiresolution analysis and pseudo-differential operators*. <http://arxiv.org/abs/0705.2294>
13. V.S. Vladimirov, I.V. Volovich and E.I. Zelenov, *p-Adic analysis and mathematical physics*. World Scientific, Singapore, 1994.

INTERNATIONAL CENTER FOR MATHEMATICAL MODELLING IN PHYSICS AND COGNITIVE SCIENCES
MSI, VÄXJÖ UNIVERSITY, SE-351 95, VÄXJÖ, SWEDEN.

E-mail address: `andrei.khrennikov@msi.vxu.se`

DEPARTMENT OF MATHEMATICS, ST.-PETERSBURG STATE ARCHITECTURE AND CIVIL ENGINEERING UNIVERSITY, 2 KRASNOARMEISKAYA 4, 190005, ST. PETERSBURG, RUSSIA.

E-mail address: `shelkv@vs1567.spb.edu`

DEPARTMENT OF APPLIED MATHEMATICS AND CONTROL PROCESSES, ST. PETERSBURG STATE UNIVERSITY, UNIVERSITetskii PR.-35, 198504 ST. PETERSBURG, RUSSIA.

E-mail address: `skopina@MS1167.spb.edu`